edénfantasys

Automatic penis pump

PRODUCT MANUAL

This product is intended for adult use as a non-medical pleasure device only. Not for medical use. No medical claims are warranted or implied by the use of this product.



CHARGING

• 3 pumping speeds

USB rechargeable

Ruler markings

Quick-release button

 Made from ABS plastic and soft TPR material

- For maximum pleasure charge before every use.
- Connect the included USB cable and insert the USB end into the wall adaptor or computer USB port (Computer must be turned on).
- The slow intermittent glow indicates charging in progress.
- A full charge takes up to 2 hours for 1.5 hours of play.

HOW TO USE

- Apply water-based lubricant on your shaft.
- Insert a flaccid or semi-flaccid penis into the donut at the base of the pump
- Hold the 😃 to turn on/off.
- Press the + button to cycle through 3 pumping speeds.
- Start pumping slowly you should feel a gentle tugging sensation.
- Once the erection is achieved, stop pumping by quickly pressing the 🙂 button.
- Release pressure by holding the button.
- If pain or discomfort occurs, immediately release pressure by holding the – button and discontinue use.

CLEANING AND CARE

- Unscrew the motorized unit and remove the donut for thorough cleaning.
- Use a toy cleaner/mild soap with water to clean the cylinder and the donut before and after each use.
- Pat with a towel and allow it to air-dry completely.
- Do not get water inside the motorized unit as it is not waterproof.

- Use with the water-based lube only, no silicone lubricants.
- Dust the donut lightly with the corn starch for better use. Store it away from heat and sunlight.

PRECAUTIONS

• Apply only the minimum amount of vacuum pressure necessary to achieve an erection; overdoing it may bruise or injure the penis.

WARNING

- Stop using the vacuum pump if pain occurs.
- Do not use under the influence of alcohol or drugs, since it may impair the user's judgment and increase the risk of injury to the penis.
- Do not use if you have sickle cell disease, have a history of prolonged erections, or you are taking large quantities of aspirin or other blood thinners, as these conditions increase the risk of bruising and hematoma.
- Use of the device may bruise or rupture the blood vessels within the penis or scrotum, resulting in petechiae, hemorrhage, or the formation of a hematoma.
- Use of the device may aggravate already existing conditions such as Peyronie's disease (the formation of hardened tissue in the penis that causes pain, curvature, and distortion, usually during erection); priapism (persistent, usually painful erection of the penis as a consequence of disease and not related to sexual arousal); and urethral strictures (urethral stricture is an area of hardened tissue which narrows the urethra, sometimes making it difficult to urinate).
- Misuse of the vacuum pump could injure the penis.
- If you have symptoms of erectile dysfunction, i.e., inability to achieve an erection that is sufficient for sexual intercourse, consult your physician prior to using this device to avoid a potentially harmful delay in diagnosing any of the most common causes of this condition, such as diabetes, multiple sclerosis, cirrhosis of the liver, chronic renal failure, or alcoholism.
- Do not use the device if you have decreased sensation of pain in the area of the penis.
- Do not use the device if you have decreased hand strength because this may make removing the device difficult.

MANAGING COMMON PROBLEMS WITH A VACUUM DEVICE

Problem	Likely cause	Remedy	Comments
No erection	No vacuum because of air leak caused by poor seal	Use more lubricant; apply firm pressure to the body; trim pubic hair; check the device for air leaks	Problem is most often associated with inexperience; it resolves as you become more adept at using the device
Partial erection	Incorrect cylinder size; incorrect insert or seal ring size; not enough practice sessions	More practice using device; use larger cylinder insert	Problem most often improves with continued use of device
Rapid loss of erection (in less than 5 minutes)	Vacuum lost because of air leak from faulty equipment, poor seal, inadequate lubrication, or excessive pubic hair	Same as for "No erection." Also: use smaller cylinder insert; use smaller or double tension rings; remove tension band carefully	Requires that you experiment with varying sizes of seal rings, inserts, and tension bands; see your physician if you need help or support to persevere
Delayed loss of erection (after 5 to 10 minutes)	Insufficient tension from bands or rings	Use smaller or combination of tension rings or bands; use the two-step application method (ask your physician about this)	
Pulling of the skin of the scrotum	Lubricant on scrotum; improper pumping technique; too large insert or seal ring; inexperience with system	Remove lubricant from scrotum; use smaller insert or seal ring; use the modified pumping technique (ask your physician about this); continue with practice sessions; angle cylinder downward during pumping	Problem may disappear as you become more adept at using the device.
Discomfort or pain:			
During pumping	Too rapid pumping; pulling of scrotal tissue	Slower pumping; continue to use device	Problem usually resolves as you become more adept at using the device
From tension ring	Tension ring too small; Anxiety	Use larger tension ring; continue to use device	Problem usually resolves as you continue to use the device
During intercourse	Inadequate lubrication; pressure on a sensitive area	More lubrication; position changes	
During ejaculation	Long period of abstinence; infection or inflammation of the prostate gland	Discontinue using device, consult your physician	
Redness, irritation, or bruising	Too rapid pumping; over pumping	Slower pumping; fewer pumps	Consult your physician if this problem persists past the first few practice sessions
Penis feels cold (to you or your partner)	Constriction of blood flow to the penis; lubrication	Wash off lubricant or warm lubricant before use	
Penis pivots on its base	Constriction of blood flow to the penis	If manageable, apply the device after you have a partial erection	



