

# Waterworks Bath Pump

## PRODUCT MANUAL

**This product is intended for adult use as a non-medical pleasure device only. Not for medical use. No medical claims are warranted or implied by the use of this product.**

- **3 pumping speeds**
- **Air-release button**
- **Ruler markings**
- **Waterproof IPX7**
- **Made from ABS plastic and soft TPE material**



## CHARGING

- For maximum pleasure, charge before every use.
- Connect the included USB cable and insert the USB end into the wall adaptor or computer USB port (Computer must be turned on).
- The slow intermittent glow indicates charging in progress.
- A full charge takes up to 150 minutes for 160 minutes of play.
- Do not use the device while charging.

## HOW TO USE

- Apply water-based lubricant onto your shaft.
- Insert a flaccid or semi-flaccid penis into the sleeve at the pump's base.
- Hold the upper  button to turn the toy on/off. The LED light will turn on.
- Press the  button to cycle through 3 pumping speeds.
- Begin pumping slowly - you should feel a gentle tugging sensation.
- Once the desired effect is achieved, stop the pumping by pressing the  button.
- Press the  button again to resume suction.
- Lightly press the  button to release the pressure gradually.
- Release pressure by holding the  button.
- If pain or discomfort occurs, immediately release pressure by holding the  button or upper  button and discontinue use.
- Lower the top outlet into the water when used in water to increase air release speed.

## CLEANING AND CARE

- Remove the sleeve or donut for thorough cleaning.
- Clean the cylinder and the donut with a toy cleaner or mild soap and water before and after each use.
- Pat with a towel and allow it to air-dry completely.
- Use with a water-based lube only, no silicone lubricants.
- To prolong its life, dust the sleeve lightly with baby powder or cornstarch. Store away from heat and sunlight.

## WARNING

- Stop using the vacuum pump if pain occurs.
- Do not use under the influence of alcohol or drugs, since it may impair the user's judgment and increase the risk of injury to the penis.
- Do not use if you have sickle cell disease, have a history of prolonged erections, or you are taking large quantities of aspirin or other blood thinners, as these conditions increase the risk of bruising and hematoma.
- Use of the device may bruise or rupture the blood vessels within the penis or scrotum, resulting in petechiae, hemorrhage, or the formation of a hematoma.
- Use of the device may aggravate already existing conditions such as Peyronie's disease (the formation of hardened tissue in the penis that causes pain, curvature, and distortion, usually during erection); priapism (persistent, usually painful erection of the penis as a consequence of disease and not related to sexual arousal); and urethral strictures (urethral stricture is an area of hardened tissue which narrows the urethra, sometimes making it difficult to urinate).
- Misuse of the vacuum pump could injure the penis.
- If you have symptoms of erectile dysfunction, i.e., inability to achieve an erection that is sufficient for sexual intercourse, consult your physician prior to using this device to avoid a potentially harmful delay in diagnosing any of the most common causes of this condition, such as diabetes, multiple sclerosis, cirrhosis of the liver, chronic renal failure, or alcoholism.
- Do not use the device if you have decreased sensation or pain in the area of the penis.
- Do not use the device if you have decreased hand strength because this may make removing the device difficult.

## MANAGING COMMON PROBLEMS WITH A VACUUM DEVICE

| Problem                                          | Likely cause                                                                                                      | Remedy                                                                                                                                                                                                      | Comments                                                                                                                                               |
|--------------------------------------------------|-------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| No erection                                      | No vacuum because of air leak caused by poor seal                                                                 | Use more lubricant; apply firm pressure to the body; trim pubic hair; check the device for air leaks                                                                                                        | Problem is most often associated with inexperience; it resolves as you become more adept at using the device                                           |
| Partial erection                                 | Incorrect cylinder size; incorrect insert or seal ring size; not enough practice sessions                         | More practice using device; use larger cylinder insert                                                                                                                                                      | Problem most often improves with continued use of device                                                                                               |
| Rapid loss of erection (in less than 5 minutes)  | Vacuum lost because of air leak from faulty equipment, poor seal, inadequate lubrication, or excessive pubic hair | Same as for "No erection." Also: use smaller cylinder insert; use smaller or double tension rings; remove tension band carefully                                                                            | Requires that you experiment with varying sizes of seal rings, inserts, and tension bands; see your physician if you need help or support to persevere |
| Delayed loss of erection (after 5 to 10 minutes) | Insufficient tension from bands or rings                                                                          | Use smaller or combination of tension rings or bands; use the two-step application method (ask your physician about this)                                                                                   |                                                                                                                                                        |
| Pulling of the skin of the scrotum               | Lubricant on scrotum; improper pumping technique; too large insert or seal ring; inexperience with system         | Remove lubricant from scrotum; use smaller insert or seal ring; use the modified pumping technique (ask your physician about this); continue with practice sessions; angle cylinder downward during pumping | Problem may disappear as you become more adept at using the device.                                                                                    |
| Discomfort or pain:                              |                                                                                                                   |                                                                                                                                                                                                             |                                                                                                                                                        |
| During pumping                                   | Too rapid pumping; pulling of scrotal tissue                                                                      | Slower pumping; continue to use device                                                                                                                                                                      | Problem usually resolves as you become more adept at using the device                                                                                  |
| From tension ring                                | Tension ring too small; Anxiety                                                                                   | Use larger tension ring; continue to use device                                                                                                                                                             | Problem usually resolves as you continue to use the device                                                                                             |
| During intercourse                               | Inadequate lubrication; pressure on a sensitive area                                                              | More lubrication; position changes                                                                                                                                                                          |                                                                                                                                                        |
| During ejaculation                               | Long period of abstinence; infection or inflammation of the prostate gland                                        | Discontinue using device, consult your physician                                                                                                                                                            |                                                                                                                                                        |
| Redness, irritation, or bruising                 | Too rapid pumping; over pumping                                                                                   | Slower pumping; fewer pumps                                                                                                                                                                                 | Consult your physician if this problem persists past the first few practice sessions                                                                   |
| Penis feels cold (to you or your partner)        | Constriction of blood flow to the penis; lubrication                                                              | Wash off lubricant or warm lubricant before use                                                                                                                                                             |                                                                                                                                                        |
| Penis pivots on its base                         | Constriction of blood flow to the penis                                                                           | If manageable, apply the device after you have a partial erection                                                                                                                                           |                                                                                                                                                        |



Customer Service 10 AM - 7 PM EST  
Phone: + 1 (770) 604 - 1352



[www.edenfantasys.com](http://www.edenfantasys.com)